



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/00, A01K 67/027, C12N 5/06	A2	(11) International Publication Number: WO 98/30683 (43) International Publication Date: 16 July 1998 (16.07.98)
(21) International Application Number: PCT/US98/00002 (22) International Filing Date: 5 January 1998 (05.01.98) (30) Priority Data: 08/781,752 10 January 1997 (10.01.97) US (71) Applicant: UNIVERSITY OF MASSACHUSETTS, A PUBLIC INSTITUTION OF HIGHER EDUCATION OF THE COMMONWEALTH OF MASSACHUSETTS [US/US]; Office of Vice Chancellor for Research at Amherst, Amherst, MA 01002 (US). (72) Inventors: STICE, Steven, L.; 468 Amherst Road, Belchertown, MA 01007 (US). CIBELLI, Jose; 166 Village Park, Amherst, MA 01002 (US). ROBL, James; 196 Old Enfield, Belchertown, MA 01007 (US). GOLUEKE, Paul; 8 Diane Drive #3, Belchertown, MA 01007 (US). PONCE DE LEON, F., Abel; 134 Wildflower Drive, Amherst, MA 01002 (US). JERRY, D., Joseph; W. Pelham Road, Shutesbury, MA 01702 (US). (74) Agents: TESKIN, Robin, L. et al.; Burns, Doane, Swecker & Mathis, L.L.P., P.O. Box 1404, Alexandria, VA 22313-1404 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: NUCLEAR TRANSFER WITH DIFFERENTIATED FETAL AND ADULT DONOR CELLS (57) Abstract <p>An improved method of nuclear transfer involving the transplantation of donor differentiated cell nuclei into enucleated oocytes of the same species as the donor cell is provided. The resultant nuclear transfer units are useful for multiplication of genotypes and transgenic genotypes by the production of fetuses and offspring, and for production of isogenic C1CM cells, including human isogenic embryonic or stem cells. Production of genetically engineered or transgenic mammalian embryos, fetuses and offspring is facilitated by the present method since the differentiated cell source of the donor nuclei can be genetically modified and clonally propagated.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

NUCLEAR TRANSFER WITH DIFFERENTIATED FETAL AND ADULT DONOR CELLS

1. FIELD OF THE INVENTION

The present invention relates to cloning procedures
5 in which cell nuclei derived from differentiated fetal or
adult, mammalian cells are transplanted into enucleated
mammalian oocytes of the same species as the donor nuclei.
The nuclei are reprogrammed to direct the development of
cloned embryos, which can then be transferred into
10 recipient females to produce fetuses and offspring, or
used to produce cultured inner cell mass cells (CICM).
The cloned embryos can also be combined with fertilized
embryos to produce chimeric embryos, fetuses and/or
offspring.

15

2. BACKGROUND OF THE INVENTION

Methods for deriving embryonic stem (ES) cell lines
in vitro from early preimplantation mouse embryos are well
known. (See, e.g., Evans et al., *Nature*, 29:154-156
(1981); Martin, *Proc. Natl. Acad. Sci., USA*, 78:7634-7638
20 (1981)). ES cells can be passaged in an undifferentiated
state, provided that a feeder layer of fibroblast cells

-2-

(Evans et al., *Id.*) or a differentiation inhibiting source (Smith et al., *Dev. Biol.*, 121:1-9 (1987)) is present.

ES cells have been previously reported to possess numerous applications. For example, it has been reported
5 that ES cells can be used as an *in vitro* model for differentiation, especially for the study of genes which are involved in the regulation of early development. Mouse ES cells can give rise to germline chimeras when introduced into preimplantation mouse embryos, thus
10 demonstrating their pluripotency (Bradley et al., *Nature*, 309:255-256 (1984)).

In view of their ability to transfer their genome to the next generation, ES cells have potential utility for germline manipulation of livestock animals by using ES
15 cells with or without a desired genetic modification.

Moreover, in the case of livestock animals, e.g., ungulates, nuclei from like preimplantation livestock embryos support the development of enucleated oocytes to term (Smith et al., *Biol. Reprod.*, 40:1027-1035 (1989);
20 and Keefer et al., *Biol. Reprod.*, 50:935-939 (1994)).

This is in contrast to nuclei from mouse embryos which beyond the eight-cell stage after transfer reportedly do

-3-

not support the development of enucleated oocytes (Cheong et al, *Biol. Reprod.*, 48:958 (1993)). Therefore, ES cells from livestock animals are highly desirable because they may provide a potential source of totipotent donor nuclei, 5 genetically manipulated or otherwise, for nuclear transfer procedures.

Some research groups have reported the isolation of purportedly pluripotent embryonic cell lines. For example, Notarianni et al., *J. Reprod. Fert. Suppl.*, 10 43:255-260 (1991), reports the establishment of purportedly stable, pluripotent cell lines from pig and sheep blastocysts which exhibit some morphological and growth characteristics similar to that of cells in primary cultures of inner cell masses isolated immunosurgically 15 from sheep blastocysts. Also, Notarianni et al., *J. Reprod. Fert. Suppl.*, 41:51-56 (1990) discloses maintenance and differentiation in culture of putative pluripotential embryonic cell lines from pig blastocysts. Gerfen et al., *Anim. Biotech*, 6(1):1-14 (1995) discloses 20 the isolation of embryonic cell lines from porcine blastocysts. These cells are stably maintained in mouse embryonic fibroblast feeder layers without the use of

-4-

conditioned medium, and reportedly differentiate into several different cell types during culture.

Further, Saito et al., *Roux's Arch. Dev. Biol.*, 201:134-141 (1992) reports cultured, bovine embryonic stem cell-like cell lines which survived three passages, but were lost after the fourth passage. Handyside et al., *Roux's Arch. Dev. Biol.*, 196:185-190 (1987) discloses culturing of immunosurgically isolated inner cell masses of sheep embryos under conditions which allow for the isolation of mouse ES cell lines derived from mouse ICMS. Handyside et al. reports that under such conditions, the sheep ICMS attach, spread, and develop areas of both ES cell-like and endoderm-like cells, but that after prolonged culture only endoderm-like cells are evident.

15 Recently, Cherny et al., *Theriogenology*, 41:175 (1994) reported purportedly pluripotent bovine primordial germ cell-derived cell lines maintained in long-term culture. These cells, after approximately seven days in culture, produced ES-like colonies which stained positive

20 for alkaline phosphatase (AP), exhibited the ability to form embryoid bodies, and spontaneously differentiated into at least two different cell types. These cells also

-5-

reportedly expressed mRNA for the transcription factors OCT4, OCT6 and HES1, a pattern of homeobox genes which is believed to be expressed by ES cells exclusively.

Also recently, Campbell et al., *Nature*, 380:64-68
5 (1996) reported the production of live lambs following nuclear transfer of cultured embryonic disc (ED) cells from day nine ovine embryos cultured under conditions which promote the isolation of ES cell lines in the mouse. The authors concluded that ED cells from day nine ovine
10 embryos are totipotent by nuclear transfer and that totipotency is maintained in culture.

Van Stekelenburg-Hamers et al., *Mol. Reprod. Dev.*,
40:444-454 (1995), reported the isolation and
characterization of purportedly permanent cell lines from
15 inner cell mass cells of bovine blastocysts. The authors isolated and cultured ICMs from 8 or 9 day bovine blastocysts under different conditions to determine which feeder cells and culture media are most efficient in supporting the attachment and outgrowth of bovine ICM
20 cells. They concluded that the attachment and outgrowth of cultured ICM cells is enhanced by the use of STO (mouse fibroblast) feeder cells (instead of bovine uterus

-6-

epithelial cells) and by the use of charcoal-stripped serum (rather than normal serum) to supplement the culture medium. Van Stekelenburg et al reported, however, that their cell lines resembled epithelial cells more than 5 pluripotent ICM cells.

Smith et al., WO 94/24274, published October 27, 1994, Evans et al, WO 90/03432, published April 5, 1990, and Wheeler et al, WO 94/26889, published November 24, 1994, report the isolation, selection and propagation of 10 animal stem cells which purportedly may be used to obtain transgenic animals. Evans et al. also reported the derivation of purportedly pluripotent embryonic stem cells from porcine and bovine species which assertedly are useful for the production of transgenic animals. Further, 15 Wheeler et al, WO 94/26884, published November 24, 1994, disclosed embryonic stem cells which are assertedly useful for the manufacture of chimeric and transgenic ungulates.

Thus, based on the foregoing, it is evident that many groups have attempted to produce ES cell lines, e.g., 20 because of their potential application in the production of cloned or transgenic embryos and in nuclear transplantation.

-7-

The use of ungulate inner cell mass (ICM) cells for nuclear transplantation has also been reported. For example, Collas et al., *Mol. Reprod. Dev.*, 38:264-267 (1994) discloses nuclear transplantation of bovine ICMs by 5 microinjection of the lysed donor cells into enucleated mature oocytes. Collas et al. disclosed culturing of embryos *in vitro* for seven days to produce fifteen blastocysts which, upon transferral into bovine recipients, resulted in four pregnancies and two births.

10 Also, Keefer et al., *Biol. Reprod.*, 50:935-939 (1994), disclosed the use of bovine ICM cells as donor nuclei in nuclear transfer procedures, to produce blastocysts which, upon transplantation into bovine recipients, resulted in several live offspring. Further, Sims et al., *Proc. Natl.* 15 *Acad. Sci., USA*, 90:6143-6147 (1993), disclosed the production of calves by transfer of nuclei from short-term *in vitro* cultured bovine ICM cells into enucleated mature oocytes.

The production of live lambs following nuclear 20 transfer of cultured embryonic disc cells has also been reported (Campbell et al., *Nature*, 380:64-68 (1996)). Still further, the use of bovine pluripotent embryonic

-8-

cells in nuclear transfer and the production of chimeric fetuses has been reported (Stice et al., *Biol. Reprod.*, 54:100-110 (1996); Collas et al, *Mol. Reprod. Dev.*, 38:264-267 (1994)). Collas et al demonstrated that 5 granulosa cells (adult cells) could be used in a bovine cloning procedure to produce embryos. However, there was no demonstration of development past early embryonic stages (blastocyst stage). Also, granulosa cells are not easily cultured and are only obtainable from females. 10 Collas et al did not attempt to propagate the granulosa cells in culture or try to genetically modify those cells.

While multiplications of genotypes are possible using embryonic cells as donors, there are problems with current methods. For example, by current methods, embryo cloning 15 can only be done using a limited number of embryonic donor nuclei (less than 100), or with in vitro cell lines. It is unknown whether the embryonic genome encodes a superior genotype until the cloned animal becomes an adult.

There also exist problems in the area of producing 20 transgenic mammals. By current methods, heterologous DNA is introduced into either early embryos or embryonic cell lines that differentiate into various cell types in the

-9-

fetus and eventually develop into a transgenic animal.

However, many early embryos are required to produce one transgenic animal and, thus, this procedure is very inefficient. Also, there is no simple and efficient

5 method of selecting for a transgenic embryo before going through the time and expense of putting the embryos into surrogate females. In addition, gene targeting techniques cannot be easily accomplished with early embryo transgenic procedures.

10 Embryonic stem cells in mice have enabled researchers to select for transgenic cells and perform gene targeting. This allows more genetic engineering than is possible with other transgenic techniques. However, embryonic stem cell lines and other embryonic cell lines must be maintained in
15 an undifferentiated state that requires feeder layers and/or the addition of cytokines to media. Even if these precautions are followed, these cells often undergo spontaneous differentiation and cannot be used to produce transgenic offspring by currently available methods.

20 Also, some embryonic cell lines have to be propagated in a way that is not conducive to gene targeting procedures.

-10-

Therefore, notwithstanding what has previously been reported in the literature, there exists a need for improved methods of cloning mammalian cells.

OBJECTS AND SUMMARY OF THE INVENTION

5 It is an object of the invention to provide novel and improved methods for producing cloned mammalian cells.

 It is a more specific object of the invention to provide a novel method for cloning mammalian cells which involves transplantation of the nucleus of a
10 differentiated mammalian cell into an enucleated oocyte of the same species.

 It is another object of the invention to provide a method for multiplying adult mammals having proven genetic superiority or other desirable traits.

15 It is another object of the invention to provide an improved method for producing genetically engineered or transgenic mammals (i.e., embryos, fetuses, offspring).

 It is a more specific object of the invention to provide a method for producing genetically engineered or
20 transgenic mammals by which a desired gene is inserted,

-11-

removed or modified in a differentiated mammalian cell or cell nucleus prior to use of that differentiated cell or cell nucleus for formation of a NT unit.

It is another object of the invention to provide
5 genetically engineered or transgenic mammals (i.e., embryos, fetuses, offspring) obtained by transplantation of the nucleus of a differentiated cell into an enucleated oocyte of the same species as the differentiated cell.

It is another object of the invention to provide a
10 novel method for producing mammalian CICM cells which involves transplantation of a nucleus of a differentiated cell into an enucleated oocyte of the same species as the differentiated cell.

It is another object of the invention to provide CICM
15 cells produced by transplantation of the nucleus of a differentiated mammalian cell into an enucleated oocyte of the same species as the differentiated cell.

It is a more specific object of the invention to provide a method for producing human CICM cells which
20 involves transplantation of nuclei of a human cell, e.g., a human adult cell, into an enucleated human oocyte.

-12-

It is another object of the invention to use such C1CM cells for therapy or diagnosis.

It is a specific object of the invention to use such C1CM cells, including human and ungulate C1CM cells, for 5 treatment or diagnosis of any disease wherein cell, tissue or organ transplantation is therapeutically or diagnostically beneficial. The C1CM cells may be used within the same species or across species.

It is another object of the invention to use tissues 10 derived from NT embryos, fetuses or offspring, including human and ungulate tissues, for treatment or diagnosis of any disease wherein cell, tissue or organ transplantation is therapeutically or diagnostically beneficial. The tissues may be used within the same species or across 15 species.

It is another specific object of the invention to use the C1CM cells produced according to the invention for the production of differentiated cells, tissues or organs.

It is a more specific object of the invention to use 20 the human C1CM cells produced according to the invention for the production of differentiated human cells, tissues or organs.

-13-

It is another specific object of the invention to use the CICM cells produced according to the invention *in vitro*, e.g. for study of cell differentiation and for assay purposes, e.g. for drug studies.

5 It is another object of the invention to provide improved methods of transplantation therapy, comprising the usage of isogenic or syngenic cells, tissues or organs produced from the CICM cells produced according to the invention. Such therapies include by way of example
10 treatment of diseases and injuries including Parkinson's, Huntington's, Alzheimer's, ALS, spinal cord injuries, multiple sclerosis, muscular dystrophy, diabetes, liver diseases, heart disease, cartilage replacement, burns, vascular diseases, urinary tract diseases, as well as for
15 the treatment of immune defects, bone marrow transplantation, cancer, among other diseases.

It is another object of the invention to provide genetically engineered or transgenic CICM cells produced by inserting, removing or modifying a desired gene in a
20 differentiated mammalian cell or cell nucleus prior to use of that differentiated cell or cell nucleus for formation of a NT unit.

-14-

It is another object of the invention to use the transgenic or genetically engineered CICM cells produced according to the invention for gene therapy, in particular for the treatment and/or prevention of the diseases and
5 injuries identified, *supra*.

It is another object of the invention to use the CICM cells produced according to the invention or transgenic or genetically engineered CICM cells produced according to the invention as nuclear donors for nuclear
10 transplantation.

Thus, in one aspect, the present invention provides a method for cloning a mammal (e.g., embryos, fetuses, offspring). The method comprises:

- (i) inserting a desired differentiated mammalian
15 cell or cell nucleus into an enucleated mammalian oocyte of the same species as the differentiated cell or cell nucleus, under conditions suitable for the formation of a nuclear transfer (NT) unit;
- (ii) activating the resultant nuclear transfer unit;
- 20 (iii) culturing said activated nuclear transfer unit until greater than the 2-cell developmental stage; and

-15-

(iv) transferring said cultured NT unit to a host mammal such that the NT unit develops into a fetus.

The cells, tissues and/or organs of the fetus are advantageously used in the area of cell, tissue and/or organ transplantation.

The present invention also includes a method of cloning a genetically engineered or transgenic mammal, by which a desired gene is inserted, removed or modified in the differentiated mammalian cell or cell nucleus prior to insertion of the differentiated mammalian cell or cell nucleus into the enucleated oocyte.

Also provided by the present invention are mammals obtained according to the above method, and offspring of those mammals.

The present invention is preferably used for cloning ungulates.

In another aspect, the present invention provides a method for producing CICM cells. The method comprises:

(i) inserting a desired differentiated mammalian cell or cell nucleus into an enucleated mammalian oocyte of the same species as the differentiated cell or cell

-16-

nucleus, under conditions suitable for the formation of a nuclear transfer (NT) unit;

(ii) activating the resultant nuclear transfer unit;

(iii) culturing said activated nuclear transfer unit
5 until greater than the 2-cell developmental stage; and

(iv) culturing cells obtained from said cultured NT unit to obtain CICM cells.

The CICM cells are advantageously used in the area of cell, tissue and organ transplantation.

10 With the foregoing and other objects, advantages and features of the invention that will become hereinafter apparent, the nature of the invention may be more clearly understood by reference to the following detailed description of the preferred embodiments of the invention
15 and to the appended claims.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides improved procedures for cloning mammals by nuclear transfer or nuclear transplantation. In the subject application, nuclear
20 transfer or nuclear transplantation or NT are used interchangeably.

-17-

According to the invention, cell nuclei derived from differentiated fetal or adult, mammalian cells are transplanted into enucleated mammalian oocytes of the same species as the donor nuclei. The nuclei are reprogrammed
5 to direct the development of cloned embryos, which can then be transferred into recipient females to produce fetuses and offspring, or used to produce CICM cells. The cloned embryos can also be combined with fertilized
10 offspring.

Prior art methods have used embryonic cell types in cloning procedures. This includes work by Campbell et al (Nature, 380:64-68, 1996) and Stice et al (Biol. Reprod., 54:100-110, 1996). In both of those studies, embryonic
15 cell lines were derived from embryos of less than 10 days of gestation. In both studies, the cells were maintained on a feeder layer to prevent overt differentiation of the donor cell to be used in the cloning procedure. The present invention uses differentiated cells.

20 It was unexpected that cloned embryos with differentiated donor nuclei could develop to advanced embryonic and fetal stages. The scientific dogma has been

-18-

that only embryonic or undifferentiated cell types could direct this type of development. It was unexpected that a large number of cloned embryos could be produced from these differentiated cell types. Also, the fact that new 5 transgenic embryonic cell lines could be readily derived from transgenic cloned embryos was unexpected.

Thus, according to the present invention, multiplication of superior genotypes of mammals, including ungulates, is possible. This will allow the 10 multiplication of adult animals with proven genetic superiority or other desirable traits. Progress will be accelerated, for example, in many important ungulate species. By the present invention, there are potentially billions of fetal or adult cells that can be harvested and 15 used in the cloning procedure. This will potentially result in many identical offspring in a short period.

The present invention also allows simplification of transgenic procedures by working with a differentiated cell source that can be clonally propagated. This 20 eliminates the need to maintain the cells in an undifferentiated state, thus, genetic modifications, both random integration and gene targeting, are more easily

-19-

accomplished. Also by combining nuclear transfer with the ability to modify and select for these cells in vitro, this procedure is more efficient than previous transgenic embryo techniques. According to the present invention, 5 these cells can be clonally propagated without cytokines, conditioned media and/or feeder layers, further simplifying and facilitating the transgenic procedure. When transfected cells are used in cloning procedures according to the invention, transgenic embryos are 10 produced which can develop into fetuses and offspring. Also, these transgenic cloned embryos can be used to produce CICM cell lines or other embryonic cell lines. Therefore, the present invention eliminates the need to derive and maintain in vitro an undifferentiated cell line 15 that is conducive to genetic engineering techniques.

The present invention can also be used to produce CICM cells, fetuses or offspring which can be used, for example, in cell, tissue and organ transplantation. By taking a fetal or adult cell from an animal and using it 20 in the cloning procedure a variety of cells, tissues and possibly organs can be obtained from cloned fetuses as they develop through organogenesis. Cells, tissues, and

-20-

organs can be isolated from cloned offspring as well.
This process can provide a source of "materials" for many
medical and veterinary therapies including cell and gene
therapy. If the cells are transferred back into the
5 animal in which the cells were derived, then immunological
rejection is averted. Also, because many cell types can
be isolated from these clones, other methodologies such as
hematopoietic chimerism can be used to avoid immunological
rejection among animals of the same species as well as
10 between species.

Thus, in one aspect, the present invention provides a
method for cloning a mammal. In general, the mammal will
be produced by a nuclear transfer process comprising the
following steps:

- 15 (i) obtaining desired differentiated mammalian cells
to be used as a source of donor nuclei;
- (ii) obtaining oocytes from a mammal of the same
species as the cells which are the source of donor nuclei;
- (iii) enucleating said oocytes;
- 20 (iv) transferring the desired differentiated cell or
cell nucleus into the enucleated oocyte, e.g., by fusion
or injection, to form NT units;

-21-

- (v) activating the resultant NT unit;
 - (vi) culturing said activated nuclear transfer unit until greater than the 2-cell developmental stage; and
 - (vii) transferring said cultured NT unit to a host
- 5 mammal such that the NT unit develops into a fetus.

The present invention also includes a method of cloning a genetically engineered or transgenic mammal, by which a desired gene is inserted, removed or modified in the differentiated mammalian cell or cell nucleus prior to

10 insertion of the differentiated mammalian cell or cell nucleus into the enucleated oocyte.

Also provided by the present invention are mammals obtained according to the above method, and offspring of those mammals. The present invention is preferably used

15 for cloning ungulates.

The present invention further provides for the use of NT fetuses and NT and chimeric offspring in the area of cell, tissue and organ transplantation.

In another aspect, the present invention provides a

20 method for producing CICM cells. The method comprises:

- (i) inserting a desired differentiated mammalian cell or cell nucleus into an enucleated mammalian oocyte

-22-

of the same species as the differentiated cell or cell nucleus, under conditions suitable for the formation of a nuclear transfer (NT) unit;

(ii) activating the resultant nuclear transfer unit;

5 (iii) culturing said activated nuclear transfer unit until greater than the 2-cell developmental stage; and

(iv) culturing cells obtained from said cultured NT unit to obtain CICM cells.

The CICM cells are advantageously used in the area of
10 cell, tissue and organ transplantation, or in the production of fetuses or offspring, including transgenic fetuses or offspring.

Preferably, the NT units will be cultured to a size of at least 2 to 400 cells, preferably 4 to 128 cells, and
15 most preferably to a size of at least about 50 cells.

Nuclear transfer techniques or nuclear transplantation techniques are known in the literature and are described in many of the references cited in the Background of the Invention. See, in particular, Campbell
20 et al, *Theriogenology*, 43:181 (1995); Collas et al, *Mol. Report Dev.*, 38:264-267 (1994); Keefer et al, *Biol. Reprod.*, 50:935-939 (1994); Sims et al, *Proc. Natl. Acad.*

-23-

Sci., USA, 90:6143-6147 (1993); WO 94/26884; WO 94/24274, and WO 90/03432, which are incorporated by reference in their entirety herein. Also, U.S. Patent Nos. 4,944,384 and 5,057,420 describe procedures for bovine nuclear
5 transplantation.

Differentiated mammalian cells are those cells which are past the early embryonic stage. More particularly, the differentiated cells are those from at least past the embryonic disc stage (day 10 of bovine embryogenesis).
10 The differentiated cells may be derived from ectoderm, mesoderm or endoderm.

Mammalian cells, including human cells, may be obtained by well known methods. Mammalian cells useful in the present invention include, by way of example,
15 epithelial cells, neural cells, epidermal cells, keratinocytes, hematopoietic cells, melanocytes, chondrocytes, lymphocytes (B and T lymphocytes), erythrocytes, macrophages, monocytes, mononuclear cells, fibroblasts, cardiac muscle cells, and other muscle cells,
20 etc. Moreover, the mammalian cells used for nuclear transfer may be obtained from different organs, e.g., skin, lung, pancreas, liver, stomach, intestine, heart,

-24-

reproductive organs, bladder, kidney, urethra and other urinary organs, etc. These are just examples of suitable donor cells. Suitable donor cells, i.e., cells useful in the subject invention, may be obtained from any cell or
5 organ of the body. This includes all somatic or germ cells.

Fibroblast cells are an ideal cell type because they can be obtained from developing fetuses and adult animals in large quantities. Fibroblast cells are differentiated
10 somewhat and, thus, were previously considered a poor cell type to use in cloning procedures. Importantly, these cells can be easily propagated in vitro with a rapid doubling time and can be clonally propagated for use in gene targeting procedures. Again the present invention is
15 novel because differentiated cell types are used. The present invention is advantageous because the cells can be easily propagated, genetically modified and selected in vitro.

Suitable mammalian sources for oocytes include sheep,
20 cows, pigs, horses, rabbits, guinea pigs, mice, hamsters, rats, primates, etc. Preferably, the oocytes will be obtained from ungulates, and most preferably bovine.

-25-

Methods for isolation of oocytes are well known in the art. Essentially, this will comprise isolating oocytes from the ovaries or reproductive tract of a mammal, e.g., a bovine. A readily available source of 5 bovine oocytes is slaughterhouse materials.

For the successful use of techniques such as genetic engineering, nuclear transfer and cloning, oocytes must generally be matured *in vitro* before these cells may be used as recipient cells for nuclear transfer, and before 10 they can be fertilized by the sperm cell to develop into an embryo. This process generally requires collecting immature (prophase I) oocytes from mammalian ovaries, e.g., bovine ovaries obtained at a slaughterhouse, and maturing the oocytes in a maturation medium prior to 15 fertilization or enucleation until the oocyte attains the metaphase II stage, which in the case of bovine oocytes generally occurs about 18-24 hours post-aspiration. For purposes of the present invention, this period of time is known as the "maturation period." As used herein for 20 calculation of time periods, "aspiration" refers to aspiration of the immature oocyte from ovarian follicles.

-26-

Additionally, metaphase II stage oocytes, which have been matured *in vivo* have been successfully used in nuclear transfer techniques. Essentially, mature metaphase II oocytes are collected surgically from either 5 non-superovulated or superovulated cows or heifers 35 to 48 hours past the onset of estrus or past the injection of human chorionic gonadotropin (hCG) or similar hormone.

The stage of maturation of the oocyte at enucleation and nuclear transfer has been reported to be significant 10 to the success of NT methods. (See e.g., Prather et al., *Differentiation*, 48, 1-8, 1991). In general, successful mammalian embryo cloning practices use the metaphase II stage oocyte as the recipient oocyte because at this stage it is believed that the oocyte can be or is sufficiently 15 "activated" to treat the introduced nucleus as it does a fertilizing sperm. In domestic animals, and especially cattle, the oocyte activation period generally ranges from about 16-52 hours, preferably about 28-42 hours post-aspiration.

20 For example, immature oocytes may be washed in HEPES buffered hamster embryo culture medium (HECM) as described in Seshagine et al., *Biol. Reprod.*, 40, 544-606, 1989, and

-27-

then placed into drops of maturation medium consisting of 50 microliters of tissue culture medium (TCM) 199 containing 10% fetal calf serum which contains appropriate gonadotropins such as luteinizing hormone (LH) and 5 follicle stimulating hormone (FSH), and estradiol under a layer of lightweight paraffin or silicon at 39°C.

After a fixed time maturation period, which ranges from about 10 to 40 hours, and preferably about 16-18 hours, the oocytes will be enucleated. Prior to 10 enucleation the oocytes will preferably be removed and placed in HECM containing 1 milligram per milliliter of hyaluronidase prior to removal of cumulus cells. This may be effected by repeated pipetting through very fine bore pipettes or by vortexing briefly. The stripped oocytes 15 are then screened for polar bodies, and the selected metaphase II oocytes, as determined by the presence of polar bodies, are then used for nuclear transfer. Enucleation follows.

Enucleation may be effected by known methods, such as 20 described in U.S. Patent No. 4,994,384 which is incorporated by reference herein. For example, metaphase II oocytes are either placed in HECM, optionally

-28-

containing 7.5 micrograms per milliliter cytochalasin B, for immediate enucleation, or may be placed in a suitable medium, for example an embryo culture medium such as CR1aa, plus 10% estrus cow serum, and then enucleated 5 later, preferably not more than 24 hours later, and more preferably 16-18 hours later.

Enucleation may be accomplished microsurgically using a micropipette to remove the polar body and the adjacent cytoplasm. The oocytes may then be screened to identify 10 those of which have been successfully enucleated. This screening may be effected by staining the oocytes with 1 microgram per milliliter 33342 Hoechst dye in HECM, and then viewing the oocytes under ultraviolet irradiation for less than 10 seconds. The oocytes that have been 15 successfully enucleated can then be placed in a suitable culture medium, e.g., CR1aa plus 10% serum.

In the present invention, the recipient oocytes will preferably be enucleated at a time ranging from about 10 hours to about 40 hours after the initiation of *in vitro* 20 maturation, more preferably from about 16 hours to about 24 hours after initiation of *in vitro* maturation, and most

-29-

preferably about 16-18 hours after initiation of *in vitro* maturation.

A single mammalian cell of the same species as the enucleated oocyte will then be transferred into the perivitelline space of the enucleated oocyte used to produce the NT unit. The mammalian cell and the enucleated oocyte will be used to produce NT units according to methods known in the art. For example, the cells may be fused by electrofusion. Electrofusion is accomplished by providing a pulse of electricity that is sufficient to cause a transient breakdown of the plasma membrane. This breakdown of the plasma membrane is very short because the membrane reforms rapidly. Thus, if two adjacent membranes are induced to breakdown and upon reformation the lipid bilayers intermingle, small channels will open between the two cells. Due to the thermodynamic instability of such a small opening, it enlarges until the two cells become one. Reference is made to U.S. Patent 4,997,384 by Prather et al., (incorporated by reference in its entirety herein) for a further discussion of this process. A variety of electrofusion media can be used including e.g., sucrose, mannitol, sorbitol and phosphate

-30-

buffered solution. Fusion can also be accomplished using Sendai virus as a fusogenic agent (Graham, Wister Inot. *Symp. Monogr.*, 9, 19, 1969).

Also, in some cases (e.g. with small donor nuclei) it
5 may be preferable to inject the nucleus directly into the oocyte rather than using electroporation fusion. Such techniques are disclosed in Collas and Barnes, *Mol. Reprod. Dev.*, 38:264-267 (1994), incorporated by reference in its entirety herein.

10 Preferably, the mammalian cell and oocyte are electrofused in a 500 μm chamber by application of an electrical pulse of 90-120V for about 15 μsec , about 24 hours after initiation of oocyte maturation. After fusion, the resultant fused NT units are then placed in a
15 suitable medium until activation, e.g., CR1aa medium. Typically activation will be effected shortly thereafter, typically less than 24 hours later, and preferably about 4-9 hours later.

The NT unit may be activated by known methods. Such
20 methods include, e.g., culturing the NT unit at sub-physiological temperature, in essence by applying a cold, or actually cool temperature shock to the NT unit. This

-31-

may be most conveniently done by culturing the NT unit at room temperature, which is cold relative to the physiological temperature conditions to which embryos are normally exposed.

5 Alternatively, activation may be achieved by application of known activation agents. For example, penetration of oocytes by sperm during fertilization has been shown to activate prefusion oocytes to yield greater numbers of viable pregnancies and multiple genetically
10 identical calves after nuclear transfer. Also, treatments such as electrical and chemical shock may be used to activate NT embryos after fusion. Suitable oocyte activation methods are the subject of U.S. Patent No. 5,496,720, to Susko-Parrish et al., herein incorporated by
15 reference in its entirety.

Additionally, activation may be effected by simultaneously or sequentially:

- (i) increasing levels of divalent cations in the oocyte, and
- 20 (ii) reducing phosphorylation of cellular proteins in the oocyte.

-32-

This will generally be effected by introducing divalent cations into the oocyte cytoplasm, e.g., magnesium, strontium, barium or calcium, e.g., in the form of an ionophore. Other methods of increasing divalent cation levels include the use of electric shock, treatment with ethanol and treatment with caged chelators.

Phosphorylation may be reduced by known methods, e.g., by the addition of kinase inhibitors, e.g., serine-threonine kinase inhibitors, such as 6-dimethyl-aminopurine, staurosporine, 2-aminopurine, and sphingosine.

Alternatively, phosphorylation of cellular proteins may be inhibited by introduction of a phosphatase into the oocyte, e.g., phosphatase 2A and phosphatase 2B.

15 In one embodiment, NT activation is effected by briefly exposing the fused NT unit to a TL-HEPES medium containing 5 μ M ionomycin and 1 mg/ml BSA, followed by washing in TL-HEPES containing 30 mg/ml BSA within about 24 hours after fusion, and preferably about 4 to 9 hours 20 after fusion.

The activated NT units may then be cultured in a suitable *in vitro* culture medium until the generation of

-33-

CICM cells and cell colonies. Culture media suitable for culturing and maturation of embryos are well known in the art. Examples of known media, which may be used for bovine embryo culture and maintenance, include Ham's F-10 + 10% fetal calf serum (FCS), Tissue Culture Medium-199 (TCM-199) + 10% fetal calf serum, Tyrodes-Albumin-Lactate-Pyruvate (TALP), Dulbecco's Phosphate Buffered Saline (PBS), Eagle's and Whitten's media. One of the most common media used for the collection and maturation of oocytes is TCM-199, and 1 to 20% serum supplement including fetal calf serum, newborn serum, estrual cow serum, lamb serum or steer serum. A preferred maintenance medium includes TCM-199 with Earl salts, 10% fetal calf serum, 0.2 mM Na pyruvate and 50 μ g/ml gentamicin sulphate. Any of the above may also involve co-culture with a variety of cell types such as granulosa cells, oviduct cells, BRL cells and uterine cells and STO cells.

Another maintenance medium is described in U.S. Patent 5,096,822 to Rosenkrans, Jr. et al., which is incorporated herein by reference. This embryo medium, named CR1, contains the nutritional substances necessary to support an embryo.

-34-

CR1 contains hemicalcium L-lactate in amounts ranging from 1.0 mM to 10 mM, preferably 1.0 mM to 5.0 mM. Hemicalcium L-lactate is L-lactate with a hemicalcium salt incorporated thereon. Hemicalcium L-lactate is significant in that a single component satisfies two major requirements in the culture medium: (i) the calcium requirement necessary for compaction and cytoskeleton arrangement; and (ii) the lactate requirement necessary for metabolism and electron transport. Hemicalcium L-lactate also serves as valuable mineral and energy source for the medium necessary for viability of the embryos.

Advantageously, CR1 medium does not contain serum, such as fetal calf serum, and does not require the use of a co-culture of animal cells or other biological media, i.e., media comprising animal cells such as oviductal cells. Biological media can sometimes be disadvantageous in that they may contain microorganisms or trace factors which may be harmful to the embryos and which are difficult to detect, characterize and eliminate.

Examples of the main components in CR1 medium include hemicalcium L-lactate, sodium chloride, potassium

-35-

chloride, sodium bicarbonate and a minor amount of fatty-acid free bovine serum albumin (Sigma A-6003).

Additionally, a defined quantity of essential and non-essential amino acids may be added to the medium. CR1
5 with amino acids is known by the abbreviation "CR1aa."

CR1 medium preferably contains the following components in the following quantities:

	sodium chloride	- 114.7 mM
	potassium chloride	- 3.1 mM
10	sodium bicarbonate	- 26.2 mM
	hemicalcium L-lactate	- 5 mM
	fatty-acid free BSA	- 3 mg/ml

In one embodiment, the activated NT embryos unit are placed in CR1aa medium containing 1.9 mM DMAP for about 4
15 hours followed by a wash in HECM and then cultured in CR1aa containing BSA.

For example, the activated NT units may be transferred to CR1aa culture medium containing 2.0 mM DMAP (Sigma) and cultured under ambient conditions, e.g., about
20 38.5°C, 5% CO₂ for a suitable time, e.g., about 4 to 5 hours.

-36-

Afterward, the cultured NT unit or units are preferably washed and then placed in a suitable media, e.g., CR1aa medium containing 10% FCS and 6 mg/ml contained in well plates which preferably contain a 5 suitable confluent feeder layer. Suitable feeder layers include, by way of example, fibroblasts and epithelial cells, e.g., fibroblasts and uterine epithelial cells derived from ungulates, chicken fibroblasts, murine (e.g., mouse or rat) fibroblasts, STO and SI-m220 feeder cell 10 lines, and BRL cells.

In one embodiment, the feeder cells comprise mouse embryonic fibroblasts. Preparation of a suitable fibroblast feeder layer is described in the example which follows and is well within the skill of the ordinary 15 artisan.

The NT units are cultured on the feeder layer until the NT units reach a size suitable for transferring to a recipient female, or for obtaining cells which may be used to produce CICM cells or cell colonies. Preferably, these 20 NT units will be cultured until at least about 2 to 400 cells, more preferably about 4 to 128 cells, and most preferably at least about 50 cells. The culturing will be

-37-

effected under suitable conditions, i.e., about 38.5°C and 5% CO₂, with the culture medium changed in order to optimize growth typically about every 2-5 days, preferably about every 3 days.

5 The methods for embryo transfer and recipient animal management in the present invention are standard procedures used in the embryo transfer industry. Synchronous transfers are important for success of the present invention, i.e., the stage of the NT embryo is in
10 synchrony with the estrus cycle of the recipient female. This advantage and how to maintain recipients are reviewed in Siedel, G.E., Jr. ("Critical review of embryo transfer procedures with cattle" in Fertilization and Embryonic Development in Vitro (1981) L. Mastroianni, Jr. and J.D.
15 Biggers, ed., Plenum Press, New York, NY, page 323), the contents of which are hereby incorporated by reference.

The present invention can also be used to clone genetically engineered or transgenic mammals. As explained above, the present invention is advantageous in
20 that transgenic procedures can be simplified by working with a differentiated cell source that can be clonally propagated. In particular, the differentiated cells used

-38-

for donor nuclei have a desired gene inserted, removed or modified. Those genetically altered, differentiated cells are then used for nuclear transplantation with enucleated oocytes.

5 Any known method for inserting, deleting or modifying a desired gene from a mammalian cell may be used for altering the differentiated cell to be used as the nuclear donor. These procedures may remove all or part of a gene, and the gene may be heterologous. Included is the
10 technique of homologous recombination, which allows the insertion, deletion or modification of a gene or genes at a specific site or sites in the cell genome.

The present invention can thus be used to provide adult mammals with desired genotypes. Multiplication of
15 adult ungulates with proven genetic superiority or other desirable traits is particularly useful, including transgenic or genetically engineered animals, and chimeric animals. Furthermore, cell and tissues from the NT fetus, including transgenic and/or chimeric fetuses, can be used
20 in cell, tissue and organ transplantation for the treatment of numerous diseases as described below in connection with the use of CICM cells.

-39-

For production of CICM cells and cell lines, after NT units of the desired size are obtained, the cells are mechanically removed from the zone and are then used.

This is preferably effected by taking the clump of cells 5 which comprise the NT unit, which typically will contain at least about 50 cells, washing such cells, and plating the cells onto a feeder layer, e.g., irradiated fibroblast cells. Typically, the cells used to obtain the stem cells or cell colonies will be obtained from the inner most

10 portion of the cultured NT unit which is preferably at least 50 cells in size. However, NT units of smaller or greater cell numbers as well as cells from other portions of the NT unit may also be used to obtain ES cells and cell colonies. The cells are maintained in the feeder 15 layer in a suitable growth medium, e.g., alpha MEM supplemented with 10% FCS and

0.1 mM β -mercaptoethanol (Sigma) and L-glutamine. The growth medium is changed as often as necessary to optimize growth, e.g., about every 2-3 days.

20 This culturing process results in the formation of CICM cells or cell lines. One skilled in the art can vary the culturing conditions as desired to optimize growth of

-40-

the particular C1CM cells. Also, genetically engineered or transgenic mammalian C1CM cells may be produced according to the present invention. That is, the methods described above can be used to produce NT units in which a
5 desired gene or genes have been introduced, or from which all or part of an endogenous gene or genes have been removed or modified. Those genetically engineered or transgenic NT units can then be used to produce genetically engineered or transgenic C1CM cells, including
10 human cells.

The resultant C1CM cells and cell lines, preferably human C1CM cells and cell lines, have numerous therapeutic and diagnostic applications. Most especially, such C1CM cells may be used for cell transplantation therapies.
15 Human C1CM cells have application in the treatment of numerous disease conditions. Human NT units *per se* may also be used in the treatment of disease conditions.

In this regard, it is known that mouse embryonic stem (ES) cells are capable of differentiating into almost any
20 cell type, e.g., hematopoietic stem cells. Therefore, human C1CM cells produced according to the invention should possess similar differentiation capacity. The C1CM

-41-

cells according to the invention will be induced to differentiate to obtain the desired cell types according to known methods. For example, the subject human C1CM cells may be induced to differentiate into hematopoietic 5 stem cells, muscle cells, cardiac muscle cells, liver cells, cartilage cells, epithelial cells, urinary tract cells, etc., by culturing such cells in differentiation medium and under conditions which provide for cell differentiation. Medium and methods which result in the 10 differentiation of C1CM cells are known in the art as are suitable culturing conditions.

For example, Palacios et al, *Proc. Natl. Acad. Sci., USA*, 92:7530-7537 (1995) teaches the production of hematopoietic stem cells from an embryonic cell line by 15 subjecting stem cells to an induction procedure comprising initially culturing aggregates of such cells in a suspension culture medium lacking retinoic acid followed by culturing in the same medium containing retinoic acid, followed by transferral of cell aggregates to a substrate 20 which provides for cell attachment.

Moreover, Pedersen, *J. Reprod. Fertil. Dev.*, 6:543-552 (1994) is a review article which references numerous

-42-

articles disclosing methods for *in vitro* differentiation of embryonic stem cells to produce various differentiated cell types including hematopoietic cells, muscle, cardiac muscle, nerve cells, among others.

5 Further, Bain et al, *Dev. Biol.*, 168:342-357 (1995) teaches *in vitro* differentiation of embryonic stem cells to produce neural cells which possess neuronal properties. These references are exemplary of reported methods for obtaining differentiated cells from embryonic or stem
10 cells. These references and in particular the disclosures therein relating to methods for differentiating embryonic stem cells are incorporated by reference in their entirety herein.

Thus, using known methods and culture medium, one
15 skilled in the art may culture the subject C1CM cells, including genetically engineered or transgenic C1CM cells, to obtain desired differentiated cell types, e.g., neural cells, muscle cells, hematopoietic cells, etc.

The subject C1CM cells may be used to obtain any
20 desired differentiated cell type. Therapeutic usages of such differentiated human cells are unparalleled. For example, human hematopoietic stem cells may be used in

-43-

medical treatments requiring bone marrow transplantation. Such procedures are used to treat many diseases, e.g., late stage cancers such as ovarian cancer and leukemia, as well as diseases that compromise the immune system, such as AIDS. Hematopoietic stem cells can be obtained, e.g., by fusing adult somatic cells of a cancer or AIDS patient, e.g., epithelial cells or lymphocytes with an enucleated oocyte, obtaining CICM cells as described above, and culturing such cells under conditions which favor differentiation, until hematopoietic stem cells are obtained. Such hematopoietic cells may be used in the treatment of diseases including cancer and AIDS.

Alternatively, adult somatic cells from a patient with a neurological disorder may be fused with an enucleated oocyte, human CICM cells obtained therefrom, and such cells cultured under differentiation conditions to produce neural cell lines. Specific diseases treatable by transplantation of such human neural cells include, by way of example, Parkinson's disease, Alzheimer's disease, ALS and cerebral palsy, among others. In the specific case of Parkinson's disease, it has been demonstrated that transplanted fetal brain neural cells make the proper

-44-

connections with surrounding cells and produce dopamine. This can result in long-term reversal of Parkinson's disease symptoms.

The great advantage of the subject invention is that
5 it provides an essentially limitless supply of isogenic or syngenic human cells suitable for transplantation. Therefore, it will obviate the significant problem associated with current transplantation methods, i.e., rejection of the transplanted tissue which may occur
10 because of host-vs-graft or graft-vs-host rejection. Conventionally, rejection is prevented or reduced by the administration of anti-rejection drugs such as cyclosporine. However, such drugs have significant adverse side-effects, e.g., immunosuppression,
15 carcinogenic properties, as well as being very expensive. The present invention should eliminate, or at least greatly reduce, the need for anti-rejection drugs.

Other diseases and conditions treatable by isogenic cell therapy include, by way of example, spinal cord
20 injuries, multiple sclerosis, muscular dystrophy, diabetes, liver diseases, i.e., hypercholesterolemia, heart diseases, cartilage replacement, burns, foot ulcers,

-45-

gastrointestinal diseases, vascular diseases, kidney disease, urinary tract disease, and aging related diseases and conditions.

This methodology can be used to replace defective genes, e.g., defective immune system genes, cystic fibrosis genes, or to introduce genes which result in the expression of therapeutically beneficial proteins such as growth factors, lymphokines, cytokines, enzymes, etc. For example, the gene encoding brain derived growth factor may be introduced into human C127 cells, the cells differentiated into neural cells and the cells transplanted into a Parkinson's patient to retard the loss of neural cells during such disease.

Previously, cell types transfected with BDNF varied from primary cells to immortalized cell lines, either neural or non-neural (myoblast and fibroblast) derived cells. For example, astrocytes have been transfected with BDNF gene using retroviral vectors, and the cells grafted into a rat model of Parkinson's disease (Yoshimoto et al., *Brain Research*, 691:25-36, (1995)).

This *ex vivo* therapy reduced Parkinson's-like symptoms in the rats up to 45% 32 days after transfer.

-46-

Also, the tyrosine hydroxylase gene has been placed into astrocytes with similar results (Lundberg et al., *Develop. Neurol.*, 139:39-53 (1996) and references cited therein).

However, such ex vivo systems have problems. In particular, retroviral vectors currently used are down-regulated in vivo and the transgene is only transiently expressed (review by Mulligan, *Science*, 260:926-932 (1993)). Also, such studies used primary cells, astrocytes, which have finite life span and replicate slowly. Such properties adversely affect the rate of transfection and impede selection of stably transfected cells. Moreover, it is almost impossible to propagate a large population of gene targeted primary cells to be used in homologous recombination techniques. By contrast, the difficulties associated with retroviral systems should be eliminated by the use of mammalian C127 cells.

Genes which may be introduced into the subject C127 cells include, by way of example, epidermal growth factor, basic fibroblast growth factor, glial derived neurotrophic growth factor, insulin-like growth factor (I and II), neurotrophin-3, neurotrophin-4/5, ciliary neurotrophic factor, AFT-1, cytokine genes (interleukins, interferons,

-47-

colony stimulating factors, tumor necrosis factors (alpha and beta), etc.), genes encoding therapeutic enzymes, etc.

In addition to the use of human C1CM cells in cell, tissue and organ transplantation, the present invention
5 also includes the use of non-human cells in the treatment of human diseases. Thus, C1CM cells, NT fetuses and NT and chimeric offspring (transgenic or non-transgenic) of any species may be used in the treatment of human disease conditions where cell, tissue or organ transplantation is
10 warranted. In general, C1CM cell, fetuses and offspring according to the present invention can be used within the same species (autologous, syngenic or allografts) or across species (xenografts). For example, brain cells from bovine NT fetuses may be used to treat Parkinson's
15 disease.

Also, the subject C1CM cells, preferably human cells, may be used as an in vitro model of differentiation, in particular for the study of genes which are involved in the regulation of early development. Also,
20 differentiated cell tissues and organs using the subject C1CM cells may be used in drug studies.

-48-

Further, the subject CICM cells may be used as nuclear donors for the production of other CICM cells and cell colonies.

In order to more clearly describe the subject invention, the following examples are provided.

-49-

EXAMPLE 1

Isolation of primary cultures of bovine and porcine embryonic and adult bovine fibroblast cells.

Primary cultures of bovine and porcine fibroblasts
5 were obtained from fetuses (45 days of pregnancy for cattle and 35 days for pig fetuses). The head, liver, heart and alimentary tract were aseptically removed, the fetuses minced and incubated for 30 minutes at 37°C in prewarmed trypsin EDTA solution (0.05% trypsin/0.02% EDTA;
10 GIBCO, Grand Island, NY). Fibroblast cells were plated in tissue culture dishes and cultured in alpha-MEM, medium (BioWhittaker, Walkersville, MD) supplemented with 10% fetal calf serum (FCS) (Hyclone, Logen, UT), penicillin (100 IU/ml) and streptomycin (50 µl/ml). The fibroblasts
15 were grown and maintained in a humidified atmosphere with 5% CO₂ in air at 37°C.

Adult fibroblast cells were isolated from the lung of a cow (approximately five years of age). Minced lung tissue was incubated overnight at 10°C in trypsin EDTA
20 solution (0.05% trypsin/0.02% EDTA; GIBCO, Grand Island, NY). The following day tissue and any disassociated cells were incubated for one hour at 37°C in prewarmed trypsin

-50-

EDTA solution (0.05% trypsin/0.02% EDTA; GIBCO, Grand Island, NY) and processed through three consecutive washes and trypsin incubations (one hr). Fibroblast cells were plated in tissue culture dishes and cultured in alpha-MEM 5 medium (BioWhittaker, Walkersville, MD) supplemented with 10% fetal calf serum (FCS) (Hyclone, Logen, UT), penicillin (100 IU/ml) and streptomycin (50 μ l/ml). The fibroblast cells can be isolated at virtually any time in development, ranging from approximately post embryonic 10 disc stage through adult life of the animal (bovine day 12 to 15 after fertilization to 10 to 15 years of age animals). This procedure can also be used to isolate fibroblasts from other mammals, including mice.

Introduction of a marker gene (foreign heterologous DNA) into embryonic and adult fibroblast cells.

The following electroporation procedure was conducted for both embryonic (cattle and pigs) and adult (cattle) fibroblast cells. Standard microinjection procedures may also be used to introduce heterologous DNA into fibroblast cells, however, in this example electroporation was used because it is an easier procedure.

-51-

Culture plates containing propagating fibroblast cells were incubated in trypsin EDTA solution (0.05% trypsin/0.02% EDTA; GIBCO, Grand Island, NY) until the cells were in a single cell suspension. The cells were spun down at 500 x g and re-suspended at 5 million cells per ml with phosphate buffered saline (PBS).

The reporter gene construct contained the cytomegalovirus promoter and the beta-galactosidase, neomycin phosphotransferase fusion gene (beta-GEO). The reporter gene and the cells at 50 µg/ml final concentration were added to the electroporation chamber. After the electroporation pulse, the fibroblast cells were transferred back into the growth medium (alpha-MEM medium (BioWhittaker, Walkersville, MD) supplemented with 10% fetal calf serum (FCS) (Hyclone, Logen, UT), penicillin (100 IU/ml) and streptomycin (50 µl/ml)).

The day after electroporation, attached fibroblast cells were selected for stable integration of the reporter gene. G418 (400 µg/ml) was added to growth medium for 15 to 20 days (range: 3 days until the end of the cultured cells' life span). This drug kills any cells without the beta-GEO gene, since they do not express the neo resistance

-52-

gene. At the end of this time, colonies of stable transgenic cells were present. Each colony was propagated independently of each other. Transgenic fibroblast cells were stained with X-gal to observe expression of beta-5 galactosidase, and confirmed positive for integration using PCR amplification of the beta-GEO gene and run out on an agarose gel.

**Use of transgenic fibroblast cells in nuclear transfer procedures to create CICM cell lines and transgenic
10 fetuses**

One line of cells (CL-1) derived from one colony of bovine embryonic fibroblast cells was used as donor nuclei in the nuclear transfer (NT) procedure. General NT procedures are described above.

15 Slaughterhouse oocytes were matured in vitro. The oocytes were stripped of cumulus cells and enucleated with a beveled micropipette at approximately 18 to 20 hrs post maturation (hpm). Enucleation was confirmed in TL-HEPES medium plus Hoechst 33342 (3 μ g/ml; Sigma). Individual
20 donor cells (fibroblasts) were then placed in the perivitelline space of the recipient oocyte. The bovine oocyte cytoplasm and the donor nucleus (NT unit) were fused together using electrofusion techniques. One fusion

-53-

pulse consisting of 120 V for 15 μ sec in a 500 μ m gap chamber was applied to the NT unit. This occurred at 24 hpm. The NT units were placed in CR1aa medium until 26 to 27 hpm.

5 The general procedure used to artificially activate oocytes has been described above. NT unit activation was initiated between 26 and 27 hpm. Briefly, NT units were exposed for four min to ionomycin (5 μ M; CalBiochem, La Jolla, CA) in TL-HEPES supplemented with 1 mg/ml BSA and
10 then washed for five min in TL-HEPES supplemented with 30 mg/ml BSA. Throughout the ionomycin treatment, NT units were also exposed to 2 mM DMAP (Sigma). Following the wash, NT units were then transferred into a microdrop of CR1aa culture medium containing 2 mM DMAP (Sigma) and
15 cultured at 38.5°C 5% CO₂ for four to five hrs. The embryos were washed and then placed in CR1aa medium plus 10% FCS and 6 mg/ml BSA in four well plates containing a confluent feeder layer of mouse embryonic fibroblast. The NT units were cultured for three more days at 38.5°C and
20 5% CO₂. Culture medium was changed every three days until days 5 to 8 after activation. At this time blastocyst stage NT embryos can be used to produce transgenic CICM

-54-

(cultured inner cell mass) cell lines or fetuses. The inner cell mass of these NT units can be isolated and plated on a feeder layer. Also, NT units were transferred into recipient females. The pregnancies were aborted at 5 35 days of gestation. This resulted in two cloned transgenic fetuses having the beta-GEO gene in all tissues checked. Thus, this is a fast and easy method of making transgenic C1CM cell lines and fetuses. This procedure is generally conducive to gene targeted C1CM cell lines and 10 fetuses.

The table below summarizes the results of these experiments.

donor cell type	n	cleavage (%)	blastocysts (%)	C1CM* lines (%)	transgenic fetuses (%)
CL-1 bovine embryonic fibroblast (bGEO)	412	220 (53)	40 (10%)	22 (55%)	
CL-1 bovine embryonic fibroblast (bGEO)	505		46 (9%)		4 fetuses†/ 16 embryos (20%)
C1CM cell line derived from CL-1 NT embryos	709		5 (0.7%)		

* 19 lines were positive for beta-GEO, 2 were negative and one line died prior to PCR detection.

-55-

† One fetus was dead and another was slightly retarded in development at 35 days of gestation. Two fetuses recovered at day 38 were normal. All fetuses were confirmed transgenic.

EXAMPLE 2

Chimeric fetuses derived from transgenic CICM cells. The transgenic CICM cell line was derived originally from a transgenic NT unit (differentiated cell).

A CICM line derived from transgenic NT embryos (a CL-1 cell transferred into an enucleated oocyte) was used to produce chimeric embryos and fetuses. Colonies of transgenic CICM cells were disaggregated either using 1-5 mg/ml pronase or 0.05% trypsin/EDTA combined with mechanical disaggregation methods so that clumps of five or fewer cells were produced. Trypsin or pronase activity was inactivated by passing the cells through multiple washes of 30 to 100% fetal calf serum. The disaggregated cells were placed in micromanipulation plates containing TL-HEPES medium. Fertilized embryos were also placed in these plates and micromanipulation tools were used to produce the chimeric embryos. Eight to ten transgenic CICM cells were injected into 8-16 cell stage fertilized embryos. These embryos were cultured in vitro to the blastocyst stage and then transferred into recipient animals.

A total of 6 blastocyst stage chimeric embryos were non-surgically transferred into two recipient females. After five weeks of gestation 3 fetuses were recovered. Several tissues of

-56-

the three fetuses, including germ cells of the gonad (suggesting germ-line chimeras), were screened by PCR amplification and southern blot hybridization of the amplified product to a beta-galactosidase fragment. Of the three fetuses, two were positive for contribution from the transgenic C127 cells. Both of these fetuses had transgenic C127 contribution to the gonad.

Transgenic NT embryos derived from transgenic C127 cell lines. The transgenic C127 cell line was derived originally from a transgenic NT unit (differentiated cell).

The same transgenic C127 cell lines were used to produce NT embryos. The NT procedures described in Example 1 were used except that C127 cells instead of fibroblast cells were used as the donor cell fused with the enucleated oocyte. Colonies of transgenic C127 cells were disaggregated either using 1-5 mg/ml pronase or 0.05% trypsin/EDTA combined with mechanical disaggregation methods so that clumps of five or fewer cells were produced. Trypsin or pronase activity was inactivated by passing the cells through multiple washes of 30 to 100% fetal calf serum before transferring the cells into enucleated oocytes. Results are reported in Table 1 (third group). Five blastocyst stage embryos were produced.

-57-

WHAT IS CLAIMED IS:

1. A method of cloning a mammal, comprising:
 - (i) inserting a desired differentiated mammalian cell or cell nucleus into an enucleated mammalian oocyte
5 of the same species as the differentiated cell or cell nucleus, under conditions suitable for the formation of a nuclear transfer (NT) unit;
 - (ii) activating the resultant nuclear transfer unit;
 - (iii) culturing said activated nuclear transfer unit
10 until greater than the 2-cell developmental stage; and
 - (iv) transferring said cultured NT unit to a host mammal such that the NT unit develops into a fetus.
2. The method according to claim 1, which further comprises developing the fetus to an offspring.
- 15 3. The method according to claim 1, wherein a desired DNA is inserted, removed or modified in said differentiated mammalian cell or cell nucleus, thereby resulting in the production of a genetically altered NT unit.

-58-

4. The method according to claim 3, which further comprises developing the fetus to an offspring.

5. The method according to claim 1, wherein the differentiated mammalian cell or cell nucleus is derived from mesoderm.

6. The method according to claim 1, wherein the differentiated mammalian cell or cell nucleus is derived from ectoderm.

7. The method according to claim 1, wherein the differentiated mammalian cell or cell nucleus is derived from endoderm.

8. The method according to claim 1, wherein the differentiated mammalian cell or cell nucleus is a fibroblast cell or cell nucleus.

9. The method according to claim 1, wherein the differentiated mammalian cell or cell nucleus is from an ungulate.

-59-

10. The method according to claim 9, wherein the ungulate is selected from the group consisting of bovine, ovine, porcine, equine, caprine and buffalo.

11. The method according to claim 1, wherein the 5 differentiated mammalian cell or cell nucleus is an adult cell or cell nucleus.

12. The method according to claim 1, wherein the differentiated mammalian cell or cell nucleus is an embryonic or fetal cell or cell nucleus.

10 13. The method according to claim 1, wherein the enucleated oocyte is matured prior to enucleation.

14. The method according to claim 1, wherein the fused nuclear transfer unit is activated by exposure to ionomycin and 6-dimethylaminopurine.

15 15. The method according to claim 3, wherein microinjection is used to insert a heterologous DNA.

-60-

16. The method according to claim 3, wherein electroporation is used to insert a heterologous DNA.

17. A fetus obtained according to the method of claim 1.

5 18. An offspring obtained according to the method of claim 2.

19. Progeny of the offspring according to claim 18.

20. A transgenic fetus obtained according to the method of claim 3.

10 21. A transgenic offspring obtained according to the method of claim 4.

22. Progeny of the offspring according to claim 21.

23. The method according to claim 1, which further comprises combining the cloned NT unit with a fertilized
15 embryo to produce a chimeric embryo.

-61-

24. The method according to claim 23, which further comprises developing the fetus to an offspring.

25. A fetus obtained according to the method of claim 23.

5 26. An offspring obtained according to the method of claim 24.

27. Progeny of the mammal according to claim 26.

28. A method of producing a CICM cell line, comprising:

- 10 (i) inserting a desired differentiated mammalian cell or cell nucleus into an enucleated mammalian oocyte of the same species as the differentiated cell or cell nucleus, under conditions suitable for the formation of a nuclear transfer (NT) unit;
- 15 (ii) activating the resultant nuclear transfer unit;
- (iii) culturing said activated nuclear transfer unit until greater than the 2-cell developmental stage; and

-62-

(iv) culturing cells obtained from said cultured NT unit to obtain a CICM cell line.

29. A CICM cell line obtained according to the method of claim 28.

5 30. The method according to claim 28, wherein a desired DNA is inserted, removed or modified in said differentiated mammalian cell or cell nucleus, thereby resulting in the production of a genetically altered NT unit.

10 31. A transgenic CICM cell line obtained according to claim 30.

32. The method of claim 28, wherein the resultant CICM cell line is induced to differentiate.

33. Differentiated cells obtained by the method of
15 claim 32.

-63-

34. Human differentiated cells obtained by the method of claim 32.

35. A method of therapy which comprises administering to a patient in need of cell transplantation therapy isogenic differentiated cells according to claim 34.

36. The method of Claim 35, wherein said cell transplantation therapy is effected to treat a disease or condition selected from the group consisting of Parkinson's disease, Huntington's disease, Alzheimer's disease, ALS, spinal cord defects or injuries, multiple sclerosis, muscular dystrophy, cystic fibrosis, liver disease, diabetes, heart disease, cartilage defects or injuries, burns, foot ulcers, vascular disease, urinary tract disease, AIDS and cancer.

37. A method of therapy which comprises administering to a human patient in need of cell transplantation therapy xenogenic differentiated cells according to claim 33.

-64-

38. The method according to claim 37 wherein the xenogenic differentiated cells are bovine cells.

39. The method of Claim 37, wherein said cell transplantation therapy is effected to treat a disease or
5 condition selected from the group consisting of Parkinson's disease, Huntington's disease, Alzheimer's disease, ALS, spinal cord defects or injuries, multiple sclerosis, muscular dystrophy, cystic fibrosis, liver disease, diabetes,
10 foot ulcers, heart disease, cartilage defects or injuries, burns, vascular disease, urinary tract disease, AIDS and cancer.

40. The method of Claim 35, wherein the differentiated human cells are hematopoietic cells or neural cells.

41. The method of Claim 35, wherein the therapy is
15 for treatment of Parkinson's disease and the differentiated cells are neural cells.

-65-

42. The method of Claim 35, wherein the therapy is for the treatment of cancer and the differentiated cells are hematopoietic cells.

43. A method of therapy which comprises
5 administering to a human patient in need of cell
transplantation therapy xenogenic cells obtained from a
fetus according to claim 17.

44. The method according to claim 43 wherein the
xenogenic cells are bovine cells.

10 45. The method of Claim 43, wherein said cell
transplantation therapy is effected to treat a disease or
condition selected from the group consisting of Parkin-
son's disease, Huntington's disease, Alzheimer's disease,
ALS, spinal cord defects or injuries, multiple sclerosis,
15 muscular dystrophy, cystic fibrosis, liver disease, diabe-
tes, heart disease, cartilage defects or injuries, burns,
foot ulcers, vascular disease, urinary tract disease, AIDS
.. and cancer.

-66-

46. A method of therapy which comprises administering to a human patient in need of cell transplantation therapy xenogenic cells obtained from an offspring according to claim 18.

5 47. The method according to claim 46 wherein the xenogenic cells are bovine cells.

48. The method of Claim 46, wherein said cell transplantation therapy is effected to treat a disease or condition selected from the group consisting of Parkinson's disease, Huntington's disease, Alzheimer's disease,
10 ALS, spinal cord defects or injuries, multiple sclerosis, muscular dystrophy, cystic fibrosis, liver disease, diabetes, heart disease, cartilage defects or injuries, burns, foot ulcers, vascular disease, urinary tract disease, AIDS
15 and cancer.

49. A method of therapy which comprises administering to a human patient in need of cell transplantation therapy xenogenic transgenic cells obtained from a transgenic fetus according to claim 20.

-67-

50. The method according to claim 49 wherein the xenogenic transgenic cells are bovine cells.

51. The method of Claim 49, wherein said cell transplantation therapy is effected to treat a disease or
5 condition selected from the group consisting of Parkinson's disease, Huntington's disease, Alzheimer's disease, ALS, spinal cord defects or injuries, multiple sclerosis, muscular dystrophy, cystic fibrosis, liver disease, diabetes, heart disease, cartilage defects or injuries, burns,
10 foot ulcers, vascular disease, urinary tract disease, AIDS and cancer.

52. A method of therapy which comprises administering to a human patient in need of cell transplantation therapy xenogenic transgenic cells
15 obtained from a transgenic offspring according to claim 21.

53. The method according to claim 52 wherein the xenogenic transgenic cells are bovine cells.

-68-

54. The method of Claim 52, wherein said cell transplantation therapy is effected to treat a disease or condition selected from the group consisting of Parkinson's disease, Huntington's disease, Alzheimer's disease, 5 ALS, spinal cord defects or injuries, multiple sclerosis, muscular dystrophy, cystic fibrosis, liver disease, diabetes, heart disease, cartilage defects or injuries, burns, foot ulcers, vascular disease, urinary tract disease, AIDS and cancer.

10 55. The method according to claim 28, which further comprises combining the cloned NT unit with a fertilized embryo to produce a chimera.

56. The method according to claim 55, which further comprises developing the chimeric CICM cell line to a 15 chimeric embryo.

57. A chimeric embryo obtained according to claim 56.

-69-

58. The method according to claim 56, which further comprises developing the chimeric embryo to a chimeric fetus.

59. A chimeric fetus obtained according to claim 58.

5 60. The method according to claim 58, which further comprises developing the chimeric fetus to a chimeric offspring.

61. A chimeric offspring obtained according to claim 60.

10 62. The method according to claim 55, wherein a desired DNA is inserted, removed or modified in said differentiated mammalian cell or cell nucleus, thereby resulting in the production of a genetically altered NT unit.

15 63. The method according to claim 62, which further comprises developing the chimeric CICM cell line to a chimeric embryo.

-70-

64. A chimeric embryo obtained according to claim
63.

65. The method according to claim 63, which further
comprises developing the chimeric embryo to a chimeric
5 fetus.

66. A chimeric fetus obtained according to claim 65.

67. The method according to claim 65, which further
comprises developing the chimeric fetus to a chimeric
offspring.

10 68. A chimeric offspring obtained according to claim
67.

69. A method of cloning a mammal, comprising:

(i) inserting a desired differentiated C1CM cell or
cell nucleus into an enucleated mammalian oocyte of the
15 same species as the differentiated C1CM cell or cell
nucleus, under conditions suitable for the formation of a
nuclear transfer (NT) unit;

-71-

(ii) activating the resultant nuclear transfer unit;
(iii) culturing said activated nuclear transfer unit
until greater than the 2-cell developmental stage; and
(iv) transferring said cultured NT unit to a host
5 mammal such that the NT unit develops into a fetus.

70. The method according to claim 69, which further
comprises developing the fetus to an offspring.

71. A fetus obtained according to the method of
claim 69.

10 72. An offspring obtained according to the method of
claim 70.

73. An organ for use as an organ xenograft, which is
obtained from the offspring according to claim 18.

74. An organ for use as an organ xenograft, which is
15 obtained from the offspring according to claim 21.

-72-

75. An organ for use as an organ xenograft, which is obtained from the offspring according to claim 26.

76. An organ for use as an organ xenograft, which is obtained from the offspring according to claim 68.

5 77. An organ for use as an organ xenograft, which is obtained from the offspring according to claim 72.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/00, A01K 67/027, C12N 5/06	A3	(11) International Publication Number: WO 98/30683 (43) International Publication Date: 16 July 1998 (16.07.98)
(21) International Application Number: PCT/US98/00002 (22) International Filing Date: 5 January 1998 (05.01.98) (30) Priority Data: 08/781,752 10 January 1997 (10.01.97) US (71) Applicant: UNIVERSITY OF MASSACHUSETTS, A PUBLIC INSTITUTION OF HIGHER EDUCATION OF THE COMMONWEALTH OF MASSACHUSETTS [US/US]; Office of Vice Chancellor for Research at Amherst, Amherst, MA 01002 (US). (72) Inventors: STICE, Steven, L.; 468 Amherst Road, Belchertown, MA 01007 (US). CIBELLI, Jose; 166 Village Park, Amherst, MA 01002 (US). ROBL, James; 196 Old Enfield, Belchertown, MA 01007 (US). GOLUEKE, Paul; 8 Diane Drive #3, Belchertown, MA 01007 (US). PONCE DE LEON, F., Abel; 134 Wildflower Drive, Amherst, MA 01002 (US). JERRY, D., Joseph; W. Pelham Road, Shutesbury, MA 01702 (US). (74) Agents: TESKIN, Robin, L. et al.; Burns, Doane, Swecker & Mathis, L.L.P., P.O. Box 1404, Alexandria, VA 22313-1404 (US).	(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 17 September 1998 (17.09.98)	
(54) Title: NUCLEAR TRANSFER WITH DIFFERENTIATED FETAL AND ADULT DONOR CELLS (57) Abstract <p>An improved method of nuclear transfer involving the transplantation of donor differentiated cell nuclei into enucleated oocytes of the same species as the donor cell is provided. The resultant nuclear transfer units are useful for multiplication of genotypes and transgenic genotypes by the production of fetuses and offspring, and for production of isogenic CICM cells, including human isogenic embryonic or stem cells. Production of genetically engineered or transgenic mammalian embryos, fetuses and offspring is facilitated by the present method since the differentiated cell source of the donor nuclei can be genetically modified and clonally propagated.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

INTERNATIONAL SEARCH REPORT

Intern Application No

PCT/US 98/00002

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/00 A01K67/027 C12N5/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A01K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CAMPBELL K H S ET AL: "SHEEP CLONED BY NUCLEAR TRANSFER FROM A CULTURED CELL LINE" NATURE, vol. 380, no. 6569, 7 March 1996, pages 64-66, XP000605296 cited in the application	1,2,9, 10,12, 13,17, 18,69-72
Y	see the whole document	28-33, 55-72
X	CAMPBELL K H S ET AL: "LIVE LAMBS BY NUCLEAR TRANSFER FROM AN ESTABLISHED CELL LINE" THERIOGENOLOGY, vol. 45, no. 1, 7 January 1996, page 287 XP000605648	1,2,9, 10,69-72
Y	see the whole document	28-33, 55-72

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

9 June 1998

Date of mailing of the international search report

07.07.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Chambonnet, F

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/00002

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 07732 A (ROSLIN INST EDINBURGH ;MCWHIR JAMES (GB); CAMPBELL KEITH HENRY STO) 14 March 1996	1-4, 9, 10, 12
Y	see page 14, line 15 - page 16, line 27	28-31, 69-72
A	<p style="text-align: center;">---</p> RITCHIE W A ET AL: "INTRACITOPLASMIC NUCLEAR INJECTION AS AN ALTERNATIVE TO CELL FUSION FOR THE PRODUCTION OF BOVINE EMBRYOS BY NUCLEAR TRANSFER" JOURNAL OF REPRODUCTION AND FERTILITY. SUPPLEMENT, vol. 5, 1 January 1995, page 60 XP000607293	1, 8-10
Y	see the whole document	28
Y	<p style="text-align: center;">---</p> COLLAS, P. & BARNES, F.L.: "Nuclear transplantation by microinjection of Inner Cell Mass and granulosa cell nuclei" MOLECULAR REPRODUCTION AND DEVELOPMENT, vol. 38, no. 3, July 1994, pages 264-267, XP002067033 cited in the application	28
A	see page 264, column 2, paragraph 2	1
Y	<p style="text-align: center;">---</p> WO 95 17500 A (ABS GLOBAL INC) 29 June 1995 see the whole document	28-31, 55-58
Y	<p style="text-align: center;">---</p> WO 95 16770 A (ABS GLOBAL INC) 22 June 1995 see the whole document	28-31, 55-70
Y	<p style="text-align: center;">---</p> CIBELLI, J.B. ET AL.: "Production of germline chimeric bovine fetuses from transgenic embryonic stem cells" THERIOGENOLOGY, vol. 47, no. 1, 17 December 1996 - January 1997, page 241 XP002067034 see the whole document & Annual conference of the international embryo transfer society, Nice, France, January 12-14 1997	28-31, 55-70
Y	<p style="text-align: center;">---</p> WO 95 34696 A (MEIJI MILK PROD CO LTD) 21 December 1995 & EP 0 774 510 A 21 May 1997 see the whole document	28-33
T	<p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	

INTERNATIONAL SEARCH REPORT

Internat al Application No

PCT/US 98/00002

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X Y	WO 97 07669 A (ROSLIN INST EDINBURGH ;CAMPBELL KEITH HENRY STOCKMAN (GB); WILMUT) 6 March 1997 see the whole document ---	1,2, 8-12,17, 18 28
P,X	WILMUT, I. ET AL.: "Viable offspring derived from fetal and adult mammalian cells" NATURE., vol. 385, no. 6619, 27 February 1997, LONDON GB, pages 810-813, XP002067035 see the whole document ---	1,2,8-12
P,X	SCHNIEKE, A.E. ET AL.: "Human Factor IX transgenic sheep produced by transfer of nuclei from transfected fetal fibroblasts" SCIENCE., vol. 278, no. 5346, 19 December 1997, LANCASTER, PA US, pages 2130-2133, XP002067036 see the whole document ---	1-4, 8-11,13, 15
P,X	WELLS, D. N. ET AL.: "Production of cloned lambs from an established embryonic cell line: a comparison between in vivo- and in vitro-matured cytoplasts" BIOLOGY OF REPRODUCTION, vol. 57, no. 2, 18 July 1997, pages 385-393, XP002067037 see the whole document ---	1,2,9,10
P,Y	WO 97 37009 A (UNIV MASSACHUSETTS A PUBLIC IN) 9 October 1997 see the whole document ---	28
P,A	WO 97 07668 A (ROSLIN INST EDINBURGH ;CAMPBELL KEITH HENRY STOCKMAN (GB); WILMUT) 6 March 1997 see the whole document ---	1
T	ZAWADA, W.M. ET AL.: "Somatic cell cloning-produced transgenic bovine neurons for transplantation in parkinsonian rats" JOURNAL OF NEUROCHEMISTRY, vol. 70, no. suppl.1, 7 March 1998, page S46 XP002067038 see the whole document ---	28-33, 35,36, 38-41

	-/--	

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/00002

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
T	<p>STICE, S.L. ET AL.: "Cloning: new breakthroughs leading to commercial opportunities" THERIOGENOLOGY, vol. 49, no. 1, January 1998, pages 129-138, XP002067444 see page 134, paragraph 2 - page 136 -----</p>	1

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 98/00002

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 1-16, 23, 24, 35-54, 63, 65, 67, 69 and 70 are directed to a method of treatment by therapy or surgery practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/00002

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9607732	A	14-03-1996	AU	3395695 A	27-03-1996
WO 9517500	A	29-06-1995	AU	1595195 A	10-07-1995
			EP	0739412 A	30-10-1996
WO 9516770	A	22-06-1995	AU	1439295 A	03-07-1995
WO 9534696	A	21-12-1995	AU	2664895 A	05-01-1996
			CA	2188944 A	21-12-1995
			EP	0759099 A	26-02-1997
			NO	965219 A	06-12-1996
			ZA	9504787 A	19-02-1996
WO 9707669	A	06-03-1997	AU	6831096 A	19-03-1997
			GB	2318578 A	29-04-1998
WO 9737009	A	09-10-1997	AU	2219497 A	22-10-1997
WO 9707668	A	06-03-1997	AU	6830996 A	19-03-1997
			GB	2318792 A	06-05-1998